

Gwent Wildlife Site/SINC Notification Form

SITE SUMMARY

SITE NAME:	Beaulieu Ridge							
GRID REF:	S	O	5	3	2	1	1	8
SIZE (approx.):	6.75ha							
DATE OF SURVEY:	2/7/18 & 3/5/19							

LOCAL AUTHORITY AREA:

Monmouthshire

WARD: Wyesham


MAIN HABITAT TYPES PRESENT:

Habitat types	NVC type	Area (ha)
Species Rich Grassland		
Marshy Grassland		

LBAP PRIORITY HABITATS/SPECIES PRESENT?

Species-rich grassland

SKETCH MAP OF SITES LOCATION:



Reproduced by permission of Ordnance Survey on behalf of HMSO. © Crown Copyright and database right 2005. All rights reserved. Ordnance Survey License number: 100016400.

FULL DESCRIPTION OF SITE (including topography, adjacent land use, access and boundaries etc.):

A 29 acre farm of which three fields (Hay Meadow, Thistle Field & Lower Field) are considered to be Local Wildlife Site quality: The hay meadow had recently been cut and the hay taken off. Although the hay field had been cut a strip of land with some bracken invasion remained uncut along the lane and fringes of uncut vegetation surrounded the rest of the field. In these uncut areas a range of meadow grasses were apparent along with coarser grasses, False Oat Grass and Cocksfoot. Other plants along these edges included Lesser Stitchwort, Common Sorrel, Hogweed. Cut-leaved Cranesbill, Red and White Clover, Lesser Trefoil, Creeping and Meadow Buttercup, Common Knapweed, Smooth Hawksbeard and an occasional plant of Yellow Rattle. Tufted Vetch was rare along the edge. The presence of these plants especially Lesser Stitchwort, Knapweed and Red Clover, indicate that the hay meadow is species-rich.

In the Thistle field there was an area to the north that was dominated by Creeping Thistles and Nettles, however generally there were numerous ant mounds and an abundance of Bird's-foot Trefoil. Perforate St John's-wort was common, also present were Knapweed, Lesser Stitchwort, Self-heal, Red Clover, Germander Speedwell, Yarrow, Meadow Buttercup and Hairy St John's-wort. Damper areas on spring lines supported Greater Birds-foot Trefoil and Marsh Bedstraw. Bracken and Bramble were encroaching along the eastern side.

The Lower Field had a series of seepages and small streams flowing down the valley sides. Some were within old Alder carr and others in more open grassland. The open seepages were dominated by Fool's Watercress and Water Mint with much Water Figwort, Brooklime Speedwell and Bog Stitchwort. In the Alder carr and along the main stream in the valley bottom, damp woodland plants predominated with species such as Wood Sorrel, Opposite-leaved Golden Saxifrage, Enchanter's Nightshade, Wood Speedwell, Ramsons, Remote Sedge and Hart's-tongue Fern. Some of the Alders along the spring-lines were mature or veteran trees. The open grassland was rich in species with abundant Bird's-foot Trefoil, Knapweed, some Bugle and Glaucous Sedge, patches of Cowslips, and much Field Woodrush, Germander Speedwell, Lesser Stitchwort and Cuckoo Flower.

Neutral Grassland:

Hay Meadow & Thistle Field = 11 indicators

Lower Field = 12 indicators

Marshy Grassland:

Lower Field = 11 indicators

Official Use Only: Site selected as a Wildlife Site/SINC? YES NO Date: 19/12/19

DETAILED SITE INFORMATION

SITE OWNERSHIP: Including names of any tenants/graziers Name(s)

Name(s)

Address(es)

[REDACTED]

[REDACTED]

Tel.no.

Mobile:

Please log all landowner/tenant contact starting from the initial permission for access to survey

Date	Contact	Discussion subjects/decisions made
[REDACTED]	[REDACTED]	[REDACTED]

Is/are the owner(s)

- i) aware of the site's nature conservation value? Yes
- ii) managing the land sympathetically? Yes
- iii) willing to manage the land sympathetically? Yes

Y	No
Y	No
Y	

Unknown

Describe the access to the site:

From Upper Beaulieu Farm

Are there any public rights of way on the site?

No

If so please describe their location:

N/A

What is the current management at the site?

Hay Field is cut for hay but was a bit early in 2018.
Other fields are grazed

Are there any threats to the site?

Cutting too soon, inappropriate tree planting, Bracken encroachment.

Are there any invasive species present?

Bracken

Are there any protected species present?

None known

Please state the section(s) of the Wildlife Sites Guidance that this site qualifies under:

H4) Neutral Grasslands - over 8 indicator species. (11 species recorded in Hay Meadow/Thistle Field and 12 in Lower Field).


Yes No

Is the site part of any grant schemes?

Grant Scheme:.....End date.....

Does the site already have any designations? Yes No Designation:.....

Sketch map of site (including northpoint, scale and areas of the site of particular note):



Please list all additional survey information both past and present that relates to this site:
CCW Phase 1, 1991.

OFFICE USE ONLY

Site no.: WS/SO51/17

Please note: The data on this form may be held in accordance with the Gwent Wildlife Trust's registration under the Data Protection Act 1998

SITE NAME: Beaulieu Ridge

FIELD NUMBER: Hay Meadow &
Thistle Field

DATE: 2/7/18

D		D	D
Agrostis capillaris (common bent)	LF	Conopodium majus (pignut)	Y
Agrostis stolonifera (creeping bent)		Crepis capillaris (smooth hawks-beard)	O
Aira praecox (early hair-grass)		Crepis biennis (rough hawks-beard)	Pteridium aquilinum (bracken)
Alopecurus geniculatus (marsh foxtail)		Dactylorhiza sp. (spotted orchid)	Pulicaria dysenterica (fleabane)
Alopecurus pratensis (meadow foxtail)		Dactylorhiza praetermissa (s.msh orchid)	Ranunculus acris (meadow buttercup)
Anthoxanthum odoratum(sweet vnl grass)	F	Daucus carota (wild carrot)	Ranunculus auricomus (gldlcks buttercup)
Arrhenatherum elatius (false oat grass)	LF	Equisetum arvense (field horsetail)	Ranunculus bulbosus (bulbous buttercup)
Brachypodium pinnatum (tor grass)		Erica cinerea (bell-heather)	Ranunculus flammula (lesser spearwort)
Briza media (quaking grass)		Erica tetralix (cross-leaved heath)	Ranunculus repens (creeping buttercup)
Bromopsis erecta (upright brome)		Eupatorium cannabinum (hemp-agrimony)	Rhinanthus minor (yellow rattle)
Bromus commutatus (meadow brome)		Euphorbia amygdaloides (wood spurge)	Rubus fruticosus (bramble)
Bromus racemosus (smooth brome)		Euphrasia officinalis agg. (eyebright)	Rumex acetosa (common sorrel)
Cynosurus cristatus (crested dog's tail)	O	Filipendula ulmaria (meadowsweet)	Rumex acetosella (Sheeps Sorrel)
Dactylis glomerata (cock's foot)		Fragaria vesca (wild strawberry)	Sanguisorba minor (salad burnet)
Danthonia decumbens (heath grass)		Galium odoratum (sweet woodruff)	Scabiosa columbaria (small scabious)
Deschampsia caespitosa (ftd hair grass)		Galium palustre (marsh-bedstraw)	Scrophularia nodosa (figwort)
Deschampsia flexuosa (wavy hair-grass)		Galium saxatile (heath bedstraw)	Senecio aquaticus (marsh ragwort)
Eriophorum angustifolium(cmn cottongrass)		Galium verum (lady's bedstraw)	Senecio jacobea (ragwort)
Festuca ovina (sheep's fescue)		Genista tinctoria (dyers greenweed)	Silaum silaus (pepper-saxifrage)
Festuca rubra (red fescue)	LF	Geranium pratense (meadow cranes bill)	Stachys officinalis (betony)
Glyceria fluitans (floating sweet-grass)		Geranium robertianum (herb robert)	Stachys palustris (marsh woundwort)
Helicotrichon pratense (mdw oat-grass)		Glechoma hederacea (ground ivy)	Stachys sylvatica (hedge woundwort)
Holcus lanatus (yorkshire fog)	F	Helianthemum nummularium (rock-rose)	Stellaria graminea (lesser stichwort)
Holcus mollis (creeping soft grass)		Heracleum sphondylium (hogweed)	O
Lolium perenne (perennial rye grass)	O	Hyacinthoides non-scripta (bluebell)	Succisa pratensis (devils bit-scabious)
Milium effusum (wood millet)		Hypericum maculatum(imp StJohns-Wort)	Taraxacum officinale (dandelion)
Melica uniflora (wood mellick)		Hypericum perforatum (pf St Johns-Wort)	O/LF
Molinia caerulea (purple moor-grass)		Hypericum tetrapterum (sq.sd StJns-Wort)	Thymus polytrichus (wild thyme)
Nardus stricta (mat grass)		Hypochaeris glabra (smooth cats ear)	Trifolium medium (zig zag clover)
Phalaris arundinacea (reed canary grass)		Hypochaeris radicata (common cats ear)	O/F
Phleum pratense (timothy grass)	O	Jasione montana (sheep's bit)	Trifolium repens (white clover)
Poa annua (annual meadow grass)		Juncus acutiflorus (sharp-flowered rush)	Tragopogon pratensis (goat's-beard)
Poa trivialis (rough meadow grass)	F	Juncus articulatus (jointed rush)	Vaccinium myrtillus (bilberry)
Trisetum flavescens (yellow oat grass)		Juncus conglomeratus (compact rush)	Valeriana dioica (marsh valerian)
Achillea millefolium (yarrow)	Y	Juncus effusus (soft rush)	Veronica beccabunga (brooklime)
Achillea ptarmica (sneezewort)		Juncus inflexus (hard rush)	Veronica chamaedrys (gemdr. speedwell)
Agrimony eupatoria (agrimony)		Juncus squarrosum (heath rush)	Vicia cracca (tufted vetch)
Ajuga reptans (bugle)		Juncus subnodulosus (blunt-flowered rush)	Vicia sativa (common vetch)
Alchemilla filicaulis (lady's mantle)		Knautia arvensis (field scabious)	Vicia sepium (bush vetch)
Alliaria petiolata (garlic mustard)		Lamiastrum galeobdolon(yellow archangel)	Viola canina (heath dog-violet)
Allium ursinum (ramsons)		Lathyrus nissolia (grass vetchling)	Viola reichenbachiana (Early dog-violet)
Anemone nemorosa (wood anemone)		Lathyrus pratensis (meadow vetchling)	Viola riviniana (common dog violet)
Angelica sylvestris (wild angelica)		Leontodon autumnalis (autumn hawkbit)	Wahlenbergia hederacea (ivy-lvd bl-flower)
Blackstonia perfoliata (yellow-wort)		Leontodon hispidus (rough hawkbit)	
Calluna vulgaris (ling heather)		Leontodon saxatilis (lesser hawkbit)	Other Species:
Caltha palustris (marsh marigold)		Leucanthemum vulgare (oxeye daisy)	<i>Bromus hordeaceus</i> (Soft Brome) R
Campanula rotundiflora (harebell)		Linum catharticum (fairy flax)	<i>Geranium dissectum</i> (Cut-leaved Cranesbill) O
Campanula trachelium (net-lvd bellflower)		Listera ovata (tway-blade orchid)	<i>Hypericum hirsutum</i> (Hairy St. Johns-wort) R
Cardamine pratensis (cuckoo flower)	Y	Lotus corniculatus (birds-foot trefoil)	<i>Rumex obtusifolius</i> (Broad-leaved Dock) Y
Carex caryophyllea (spring sedge)		Lotus uliginosus(greater birds-foot trefoil)	<i>Trofolium dubium</i> (Lesser Trefoil) O
Carex disticha (brown sedge)		Luzula campestris (field wood rush)	
Carex divisa (grey sedge)		Luzula multiflora (heath wood-rush)	
Carex obtusata (false fox sedge)		Lychis flos-cuculi (ragged robin)	
Carex flacca (glauconous sedge)		Melampyrum pratense (cow-wheat)	
Carex nigra (common sedge)		Mentha aquatica (water mint)	
Carex ovalis (oval sedge)		Mercurialis perennis (dog's-mercury)	
Carex panicea (carnation sedge)		Myosotis arvensis (field forget-me-not)	
Carex pseudocyperus (cyperus sedge)		Narcissus pseudonarcissus (wild daffodill)	
Carex remota (Remote sedge)		Oenanthe crocata (Hemlock water dropwort)	
Carex riparia (greater pond-sedge)		Ononis repens (common restarrow)	
Carex rostrata (bottle sedge)		Ophrys apifera (bee orchid)	
Carex spicata (spiked sedge)		Orchis mascula (early purple orchid)	

Carex sylvatica (wood sedge)		Orchis morio (green-winged orchid)		
Centaurea nigra (common knapweed)	O/LF	Pilosella officinarum (mse-ear hawkweed)		
Centaurea scabiosa (greater knapweed)		Pimpinella saxifraga (burnet saxifrage)		
Centaurium erythraea (cmnn centuary)		Plantago lanceolata (ribwort plantain)	F	
Cerastium holosteoides(cmn. mouse-ear)	Y	Polygala serpyllifolia (heath milkwort)		
Chrysosplenium oppositifolium (op.l.gsxfg)		Polygala vulgaris (common milkwort)		
Cirsium arvense (creeping thistle)	LA	Potentilla anserina (silverweed)		
Cirsium dissectum (meadow thistle)		Potentilla erecta (tormentil)		
Cirsium palustre (marsh thistle)	O	Potentilla reptans (creeping cinquefoil)		
Cirsium vulgare (spear thistle)	VO	Potentilla sterilis (barren strawberry)		

SITE NAME: Beaulieu Ridge

FIELD NUMBER: Lower Field

DATE: 3/5/19

D		D	D
Agrostis capillaris (common bent)		Conopodium majus (pignut)	O
Agrostis stolonifera (creeping bent)	Y	Crepis capillaris (smooth hawks-beard)	Prunella vulgaris (self-heal)
Aira praecox (early hair-grass)		Crepis biennis (rough hawks-beard)	Pteridium aquilinum (bracken)
Alopecurus geniculatus (marsh foxtail)		Dactylorhiza sp. (spotted orchid)	Pulicaria dysenterica (fleabane)
Alopecurus pratensis (meadow foxtail)	Y	Dactylorhiza praetermissa (s.msh orchid)	Ranunculus acris (meadow buttercup)
Anthoxanthum odoratum(sweet vnl grass)	Y	Daucus carota (wild carrot)	Ranunculus auricomus (gldlcks buttercup)
Arrhenatherum elatius (false oat grass)		Equisetum arvense (field horsetail)	Ranunculus bulbosus (bulbous buttercup)
Brachypodium pinnatum (tor grass)		Erica cinerea (bell-heather)	Ranunculus flammula (lesser spearwort)
Briza media (quaking grass)		Erica tetralix (cross-leaved heath)	Ranunculus repens (creeping buttercup)
Bromopsis erecta (upright brome)		Eupatorium cannabinum (hemp-agrimony)	Rhinanthus minor (yellow rattle)
Bromus commutatus (meadow brome)		Euphorbia amygdaloides (wood spurge)	Rubus fruticosus (bramble)
Bromus racemosus (smooth brome)		Euphrasia officinalis agg. (eyebright)	Rumex acetosa (common sorrel)
Cynosurus cristatus (crested dog's tail)		Filipendula ulmaria (meadowsweet)	Rumex acetosella (Sheeps Sorrel)
Dactylis glomerata (cock's foot)		Fragaria vesca (wild strawberry)	Sanguisorba minor (salad burnet)
Danthonia decumbens (heath grass)		Galium odoratum (sweet woodruff)	Scabiosa columbaria (small scabious)
Deschampsia caespitosa (ftd hair grass)		Galium palustre (marsh-bedstraw)	Scrophularia nodosa (figwort)
Deschampsia flexuosa (wavy hair-grass)		Galium saxatile (heath bedstraw)	Senecio aquaticus (marsh ragwort)
Eriophorum angustifolium(cmn cottongrass)		Galium verum (lady's bedstraw)	Senecio jacobaea (ragwort)
Festuca ovina (sheep's fescue)		Genista tinctoria (dyers greenweed)	Silaum silaus (pepper-saxifrage)
Festuca rubra (red fescue)		Geranium pratense (meadow cranes bill)	Stachys officinalis (betony)
Glyceria fluitans (floating sweet-grass)		Geranium robertianum (herb robert)	Stachys palustris (marsh woundwort)
Helictotrichon pratense (mdw oat-grass)		Glechoma hederacea (ground ivy)	Y
Holcus lanatus (yorkshire fog)		Helianthemum nummularium (rock-rose)	Stachys sylvatica (hedge woundwort)
Holcus mollis (creeping soft grass)		Heracleum sphondylium (hogweed)	Y
Lolium perenne (perennial rye grass)		Hyacinthoides non-scripta (bluebell)	Stellaria graminea (lesser stichwort)
Milium effusum (wood millet)		Hypericum maculatum(imp StJohns-Wort)	O
Melica uniflora (wood mellick)		Hypericum perforatum (pf St Johns-Wort)	Taraxacum officinale (dandelion)
Molinia caerulea (purple moor-grass)		Hypericum tetrapterum (sq.sd StJns-Wort)	Y
Nardus stricta (mat grass)		Hypochaeris glabra (smooth cats ear)	Trifolium medium (zig zag clover)
Phalaris arundinacea (reed canary grass)		Hypochaeris radicata (common cats ear)	O
Phleum pratense (timothy grass)		Jasione montana (sheep's bit)	Trifolium repens (white clover)
Poa annua (annual meadow grass)		Juncus acutiflorus (sharp-flowered rush)	Tragopogon pratensis (goat's-beard)
Poa trivialis (rough meadow grass)	Y	Juncus articulatus (jointed rush)	Vaccinium myrtillus (bilberry)
Trisetum flavescens (yellow oat grass)		Juncus conglomeratus (compact rush)	Valeriana dioica (marsh valerian)
Achillea millefolium (yarrow)	Y	Juncus effusus (soft rush)	Valeriana officinalis (common valerian)
Achillea ptarmica (sneezewort)		Juncus inflexus (hard rush)	Veronica beccabunga (brooklime)
Agrimonia eupatoria (agrimony)		Juncus squarrosum (heath rush)	Veronica chamaedrys (gemdr. speedwell)
Ajuga reptans (bugle)	Y	Juncus subnodulosus (blunt-flowered rush)	Vicia cracca (tufted vetch)
Alchemilla filicaulis (lady's mantle)		Knautia arvensis (field scabious)	Vicia sativa (common vetch)
Alliaria petiolata (garlic mustard)		Lamiastrum galeobdolon(yellow archangel)	Vicia sepium (bush vetch)
Allium ursinum (ramsons)		Lathyrus nissolia (grass vetchling)	Viola canina (heath dog-violet)
Anemone nemorosa (wood anemone)		Lathyrus pratensis (meadow vetchling)	Viola reichenbachiana (Early dog-violet)
Angelica sylvestris (wild angelica)	Y	Leontodon autumnalis (autumn hawkbit)	Viola riviniana (common dog violet)
Blackstonia perfoliata (yellow-wort)		Leontodon hispidus (rough hawkbit)	Wahlenbergia hederacea (ivy-lvd bl-flower)
Calluna vulgaris (ling heather)		Leontodon saxatilis (lesser hawkbit)	Other Species:
Caltha palustris (marsh marigold)		Leucanthemum vulgare (oxeye daisy)	<i>Anemone nemorosa</i> (Wood Anemone)
Campanula rotundiflora (harebell)		Linum catharticum (fairy flax)	<i>Anisantha sterilis</i> (Barren Brome)
Campanula trachelium (net-lvd bellflower)		Listera ovata (tway-blade orchid)	<i>Urtica dioica</i> (Stinging Nettle)
Cardamine pratensis (cuckoo flower)	Y	Lotus corniculatus (birds-foot trefoil)	Apium nodiflorum (Fools Watercress)
Carex caryophyllea (spring sedge)		Lotus uliginosus(greater birds-foot trefoil)	Anthriscus sylvestris (Cow Parsley)
Carex disticha (brown sedge)		Luzula campestris (field wood rush)	Arabidopsis thaliana (Thale Cress)
Carex divisa (grey sedge)		Luzula multiflora (heath wood-rush)	Arum maculatum (Lords & Ladies)

Carex obtrudea (false fox sedge)		<i>Lychnis flos-cuculi</i> (ragged robin)		Asplenium scolopendrium (Hart's Tongue)	Y
Carex flacca (glaucous sedge)	Y	Melampyrum pratense (cow-wheat)		Athyrium filix-femina (Lady Fern)	Y
Carex nigra (common sedge)		<i>Mentha aquatica</i> (water mint)	Y	Bellis perennis (Daisy)	Y
Carex ovalis (oval sedge)		Mercurialis perennis (dog's-mercury)	Y	Bromus hordeaceus (Soft Brome)	Y
Carex panicea (carnation sedge)		Myosotis arvensis (field forget-me-not)	Y	Capsella bursa-pastoris (Shepherd's Purse)	Y
Carex pseudocyperus (cyperus sedge)		<u>Narcissus pseudonarcissus</u> (wild daffodill)		Cardamine flexuosa (Wavy Bitter-cress)	Y
Carex remota (Remote sedge)	Y	<i>Oenanthe crocata</i> (Hemlock water dropwort).		Cardamine hirsuta (Hairy Bitter-cress)	Y
Carex riparia (greater pond-sedge)		Ononis repens (common restarrow)		Cerastium glomeratum (Sticky Mouse-ear)	Y
Carex rostrata (bottle sedge)		<u>Ophrys apifera</u> (bee orchid)		<u>Chrysosplenium oppositifolium</u> (Opposite-leaved Golden Saxifrage)	Y
Carex spicata (spiked sedge)		Orchis mascula (early purple orchid)		Circaea lutetiana (Enchanter's Nightshade)	Y
Carex sylvatica (wood sedge)		Orchis morio (green-winged orchid)		Crataegus monogyna (Hawthorn)	Y
Centaurea nigra (common knapweed)	Y	Pilosella officinarum (mse-ear hawkweed)		Digitalis purpurea (Foxglove)	Y
Centaurea scabiosa (greater knapweed)		Pimpinella saxifraga (burnet saxifrage)		Euonymus europaeus (Spindle)	Y
Centaurium erythraea (cmnn centuary)		Plantago lanceolata (ribwort plantain)	F	Euphorbia peplus (Petty Spurge)	Y
Cerastium holosteoides(cmnn. mouse-ear)	Y	Polygala serpyllifolia (heath milkwort)		Fraxinus excelsior (Ash)	Y
Chrysosplenium oppositifolium (op.l.gsxfg)		Polygala vulgaris (common milkwort)		Geranium dissectum (Cut-leaved Cranesbill)	Y
Cirsium arvense (creeping thistle)	Y	Potentilla anserina (silverweed)		Lamiastrum galeobdolon (Yellow Archangel)	Y
Cirsium dissectum (meadow thistle)		Potentilla erecta (tormentil)		Allium ursinum (Ramsone)	Y
Cirsium palustre (marsh thistle)	Y	Potentilla reptans (creeping cinquefoil)	Y	<i>Alnus glutinosa</i> (Alder)	Y
Cirsium vulgare (spear thistle)		Potentilla sterilis (barren strawberry)	Y	Lamium album (White Dead-Nettle)	Y
				Lepidium coronopus (Swine Cress)	Y
				<u>Lysimachia nemorum</u> (Yellow Pimpernel)	Y
				Moehringia trinervia (Three-nerved Sandwort)	Y
				Myosotis sylvatica (Wood Forget-me-not)	Y
				Oxalis acetosella (Wood Sorrel)	Y
				Pentaglottis sempervirens (Green Alkanet)	Y
				Rumex obtusifolius (Broad-leaved Dock)	Y
				Salix viminalis (Osier)	Y
				<u>Scrophularia auriculata</u> (Water Figwort)	Y
				Silene dioica (Red Campion)	Y
				Solanum dulcamara (Bittersweet)	Y
				Sonchus asper (Prickly Sow-thistle)	Y
				<u>Stellaria alsine</u> (Bog Stitchwort)	Y
				Stellaria holostea (Greater Stitchwort)	Y
				Urtica dioica (Stinging Nettle)	Y
				Veronica arvensis (Wall Speedwell)	Y
				Veronica hederifolia (Ivy-leaved Speedwell)	Y
				Veronica montana (Wood Speedwell)	Y
				Veronica serpyllifolia (Thyme-leaved Speedwell)	Y
				Vicia hirsute (Hairy Tare)	Y

Key to colour coding of grassland species list

Species in **Bold** equals indicator species for Neutral grasslands

Species in **Blue** equals indicator species for Marshy grasslands

(Please note other species in list may be indicator species for other grassland types and these have been taken in to account when assessing overall value of the site).

Species in **Bold and underlined** equals a species from the contributory species list.

Species in ***red italics*** equals a species from the primary species list.

D = Dominant

A = Abundant

F = Frequent

O = Occasional

R = Rare

V = Very

L = Local/Locally

Y = Present