## CYNGOR CEFN GWLAD CYMRU COUNTRYSIDE COUNCIL FOR WALES

## SITE OF SPECIAL SCIENTIFIC INTEREST CITATION

## POWYSGWAUN EFAIL-LLWYDIARTHDate of Notification:1990National Grid Reference:SJ 069169O.S. Maps:1:50,000 Sheet number: 125<br/>1:25,000 Sheet number: SJ 01Site Area:2.4 ha

## **Description:**

The site comprises a single damp pasture field near Llanfihangel-yng-Ngwynfa. It supports one of the finest stands of marsh vegetation known in Montgomeryshire, along with areas of species-rich neutral grassland.

A small brook runs down the centre of the field. Marshy ground on either side is characterised by an abundance of sharp-flowered rush Juncus acutiflorus and jointed rush J. articulatus, along with frequent sweet vernal-grass Anthoxanthum odoratum, quakinggrass Briza media, crested dog's-tail Cynosurus cristatus, common sedge Carex nigra, marsh-marigold Caltha palustris, water horsetail Equisetum fluviatile, greater bird's-foottrefoil Lotus uliginosus, creeping buttercup Ranunculus repens and the locally uncommon northern marsh-orchid Dactylorhiza purpurella. Other associated species include carnation sedge Carex panicea, marsh willowherb Epilobium palustre, wood horsetail Equisetum sylvaticum, common marsh-bedstraw Galium palustre, ragged-robin Lychnis flos-cuculi, water mint Mentha aquatica, meadow buttercup Ranunculus acris and marsh arrowgrass Triglochin palustris. This community grades into drier grassland on the banks above, dominated by common bent Agrostis capillaris, sweet vernal-grass, crested dog's tail and red fescue Festuca rubra, with frequent common knapweed *Centaurea nigra*, ribwort plantain *Plantago lanceolata*, meadow buttercup and clovers Trifolium spp., and a range of other species such as intermediate lady's-mantle Alchemilla xanthochlora, pignut Conopodium majus, autumn hawkbit Leontodon autumnalis, rough hawkbit L. hispidus, common bird's-foot-trefoil Lotus corniculatus, bulbous buttercup Ranunculus bulbosus and common sorrel Rumex acetosa. Certain banks support a variant of this community with frequent mat-grass Nardus stricta, carnation sedge, tormentil Potentilla erecta and devil's-bit scabious Succisa pratensis. Additional associates here include bugle Ajuga reptans, heath spotted-orchid Dactylorhiza maculata subsp. ericetorum, lousewort Pedicularis sylvatica and yellowrattle Rhinanthus minor. The more level areas of dry pasture have been affected by nutrient enrichment and cattle poaching – they support a modified sward type in which perennial rye-grass Lolium perenne, thistles Cirsium spp. and docks Rumex spp. are common.

This document is **NOT** a definitive legal version and has been formatted, updated and partially edited for use on the CCW Web site. This document should not be used in any legal proceedings, public enquiry or any other hearing or appeal. If you require a full legal copy of the document please contact CCW in writing.