# CYNGOR CEFN GWLAD CYMRU COUNTRYSIDE COUNCIL FOR WALES

### SITE OF SPECIAL SCIENTIFIC INTEREST CITATION

### POWYS

## **CAE LLWYN**

Date of Notification:	1990
National Grid Reference:	SO098634
<u>O.S. Maps:</u>	1:50,000 Sheet number: 147 1:25,000 Sheet number: SO06
Site Area:	3.4 ha

#### **Description:**

This site comprises a single, gently sloping field with a southerly aspect. It is an excellent example of a traditionally managed herb-rich meadow supporting both "wet" and "dry" plant communities.

The top part of the meadow supports a rather impoverished community dominated by perennial rye-grass *Lolium perenne* and crested dog's-tail *Cynosurus cristatus*, but this soon gives way to a more typical unimproved grassland community on the slopes below. Common bent *Agrostis capillaris*, crested dog's-tail and Yorkshire-fog *Holcus lanatus* are dominant here, along with frequent sweet vernal-grass *Anthoxanthum odoratum*, pignut *Conopodium majus*, common bird's-foot-trefoil *Lotus corniculatus*, ribwort plantain *Plantago canceolata* and clovers *Trifolium spp*. Other typical associates include common knapweed *Centaurea nigra*, meadow vetchling *Lathyrus pratensis*, rough hawkbit *Leontodon hispidus*, yellow-rattle *Rhinanthus minor*, common sorrel *Rumex acetosa*, great burnet *Sanguisorba officinalis* and tufted vetch *Vicia cracca*.

The lower part of the field supports a rich fen-meadow community dominated by mixtures of sweet vernal-grass, crested dog's-tail, red fescue *Festuca rubra*, Yorkshire-fog and purple moor-grass *Molinia caerulea*, with frequent carnation sedge *Carex panicea*, meadow thistle *Cirsium dissectum*, autumn hawkbit *Leontodon autumnalis*, ribwort plantain, tormentil *Potentilla erecta*, meadow buttercup *Ranunculus acris*, yellow-rattle, devil's-bit scabious *Succisa pratensis* and clovers. Other associated species include quaking-grass *Briza media*, bugle *Ajuga reptans*, common knapweed, heath-spotted orchid *Dactylorhiza maculata subsp. ericetorum*, sharp-flowered rush *Juncus acuiflorus*, bitter-vetch *Lathyrus montanus*, lousewort *Pedicularis sylvatica*, saw-wort *Serratula tinctoria*, betony *Stachys officinalis* and the locally uncommon dyer's greenweed *Genista tinctoria*. Wet hollows dominated by sharp-flowered rush support additional species such as sneezewort *Achillea ptarmica*, common spotted-orchid *Dactylorhiza fuchsii* and meadowsweet *Filipendula ulmaria*.

This document is **NOT** a definitive legal version and has been formatted, updated and partially edited for use on the CCW Web site. This document should not be used in any legal proceedings, public enquiry or any other hearing or appeal. If you require a full legal copy of the document please contact CCW in writing.